

You should read this written explanation before use. Please keep it so that you can read it when you need to.



「MORI」 ZENTI Tablets

[Category ㊗ OTC medicines]

Indication

Bronchial asthma, bronchial catarrh, pleuritis,
Pulmonary tuberculosis, common cold cough



! CAUTION FOR USE

Consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of physical strength or constitutional weakness).
 - (4) Persons who have gastrointestinal weakness.
 - (5) Persons with excessive sweating.
 - (6) Elderly persons.
 - (7) Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
 - (8) Persons with the following symptoms : Edema, urination difficulty.
 - (9) Persons diagnosed as having the following : Hypertension, heart disease, kidney disease, thyroid disease.
2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Nausea, loss of appetite, gastric distress

Information sheet for customer

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptom's name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually worsen.

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month (5 to 6 days if you take this medicine for the common cold).
4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

Dosage and administration

Take the following doses with water or warm water in the morning and evening on an empty stomach.

Age	One dosage	Daily dosage
16 years and over	3 ~ 6 tablets	2 times a day
8 to 15 years old	2 ~ 3 tablets	

〈Precautions of Dosage and Administration〉

When allowing children to take the medicine, a guardian must be present to watch and tell such them how to take it.

Ingredient and amount

6 tablets (2.25 g) of this medicine contains 1.05 g of the dried extract from the following mixed crude drugs.

Pinellia Tuber	1.5 g	Peony Root	1 g
Glycyrrhiza	1 g	Ginger	1 g
Platycodon Root	1 g	Ephedra Herb	1 g
Cinnamon Bark	1 g	Schisandra Fruit	0.5g
Asiasarum Root	1 g	Citrus Unshiu Pee	0.5g

The inactive ingredients contained are Refined White Sugar, Talc, Gum Arabic, Titanium Oxide, Magnesium Stearate, Polyvinyl Acetal Diethylaminoacetate, Carnauba Wax.

Precautions for storage and handling

- (1) Store in a cool, low humidity place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer the medicine to a different container.
(It may lead to misuse or alter the quality of the drug.)

REQUEST FOR LITERATURE SHOULD BE MADE TO :

Dep. of Medical Information,

Ohsugi Pharmaceutical Co., Ltd.

1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035

Website <https://www.ohsugi-kanpo.co.jp>