



# MORI MASHIN

## (Mashinginan Extract Granules)

[Category 2 OTC medicines]

### Indication

Following symptoms in persons with moderate or lower physical strength who have hard stools like rabbit dung in some case :

Constipation and its accessory symptoms such as heavy head, hot flashes, eczema, dermatitis, pimple (acne), loss of appetite (anorexia), abdominal bloating, abnormal intestinal fermentation and hemorrhoid.



## ! CAUTION FOR USE

### To be avoided

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

1. Don't give this medicine to children under 3 months of age.
2. This medicine should not be taken together with the following medicines :  
Other cathartic agents.
3. Women who are nursing should not be taken this medicine. If they take it, avoid breast feeding.

### Consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.
  - (1) Persons undergoing medical treatment from a physician.
  - (2) Pregnant women or women suspected of being pregnant.
  - (3) Persons who have a gastrointestinal weakness and tend to diarrhea.

## Information sheet for customer

2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Gastrointestinal system	Diarrhea with severe abdominal pain, abdominal pain

3. A following symptom may appear after taking this medicine. If the symptom continues or worsen, consult a physician, pharmacist or registered distributor with this leaflet :  
Diarrhea

4. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for 5 to 6 days.

## Dosage and administration

Take the following doses with water or warm water before or between meals.

(Between meals means 2-3 hours after meals.)

Age	One dosage	Daily dosage
15 years and over	1 packet (2.0g)	3 times a day
7 to 14 years old	2/3 packet	
4 to 6 years old	1/2packet	
2 to 3 years old	1/3 packet	
Less than 2 years old	1/4 packet	

〈Precautions of Dosage and Administration〉

- (1) When allowing children to take the medicine, a guardian must be present to watch and tell such them how to take it.
- (2) For infants younger than 1 year old, medical care by a physician should always precede administration of this drug. This drug should be administered only if it is absolutely necessary.

## Ingredient and amount

3 packets (2.0 g × 3 : 6.0 g) of this medicine contains 2.6 g of the dried extract (Mashiningan extract) from the following mixed crude drugs.

Hemp Fruit	5 g	Magnolia Bark	2 g
Peony Root	2 g	Rhubarb	4 g
Immature Orange	2 g	Apricot Kernel	2 g

The inactive ingredients contained are Lactose Hydrate, Corn Starch and Magnesium Stearate.

## **Precautions for storage and handling**

- (1) Store in a cool, low humidity place away from direct sunlight.
  - (2) Keep out of reach of children.
  - (3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
  - (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.
- This product is manufactured from natural crude drugs. There may be slight differences in the color and taste of the product. There is no change in the effect.

REQUEST FOR LITERATURE SHOULD BE MADE TO :

Dep. of Medical Information,

Ohsugi Pharmaceutical Co., Ltd.

1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035

Website <https://www.ohsugi-kanpo.co.jp>