



MORI KOMYOU

(Shichimotsukokato Extract Granules)

[Category 2 OTC medicines]

Indication

Following symptoms in persons with moderate or poor physical strength who look pale, get tired easily and have not gastrointestinal disorders :

Accessory symptoms of hypertension (hot flashes, stiff shoulder, ear noise, heavy head).



! CAUTION FOR USE

Consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

- (1) Persons undergoing medical treatment from a physician.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Persons who have a gastrointestinal weakness and tend to diarrhea.

2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress

3. A following symptom may appear after taking this medicine. If the symptom continues or worsen, consult a physician, pharmacist or registered distributor. Take this leaflet with you : Diarrhea.

4. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.

Dosage and administration

Take the following doses with water or warm water before or between meals.
(Between meals means 2-3 hours after meals.)

Age	One dosage	Daily dosage
15 years and over	1 packet (2.5g)	3 times a day
Less than 15 years old	Do not use	

Ingredient and amount

3 packets (2.5 g × 3 : 7.5 g) of this medicine contains 4.3 g of the dried extract (Shichimotsukokato extract) from the following mixed crude drugs.

Japanese Angelica Root	3 g	Uncaria Hook	4 g
Peony Root	3 g	Astragalus Root	3 g
Cnidium Rhizome	3 g	Phellodendron Bark	2 g
Rehmannia Root	3 g		

The inactive ingredients contained are Lactose Hydrate, Corn Starch and Magnesium Stearate.

Precautions for storage and handling

- (1) Store in a cool, low humidity place away from direct sunlight.
 - (2) Keep out of reach of children.
 - (3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
 - (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.
- This product is manufactured from natural crude drugs. There may be slight differences in the color and taste of the product. There is no change in the effect.

REQUEST FOR LITERATURE SHOULD BE MADE TO :

Dep. of Medical Information,

Ohsugi Pharmaceutical Co., Ltd.

1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035

Website <https://www.ohsugi-kanpo.co.jp>