



MORI HAITUUN

(Sokeikakketsuto Extract Granules)

[Category 2 OTC medicines]

Indication

Following symptoms in persons with moderate physical strength who have pain and have numbness in some case :
Joint pain, nerve pain, low back pain, and muscle pain.



! CAUTION FOR USE

To be avoided

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)

Don't give this medicine to children under 3 months of age.

Consultation

1. The following persons should consult a physician, pharmacist or registered distributor before taking this medicine.

- (1) Persons undergoing medical treatment from a physician.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Persons who have a gastrointestinal weakness and tend to diarrhea.
- (4) Elderly persons.
- (5) Persons who have experienced rash, redness, itching, etc. due to medicines, etc.
- (6) Persons with the following symptom : Edema.
- (7) Persons diagnosed as having the following : Hypertension, heart disease, kidney disease.

Information sheet for customer

2. The following symptoms may be adverse reactions of this medicine. If any of the following symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist or registered distributor with this instruction leaflet.

Areas	Symptoms
Skin	Rash, redness, itching
Gastrointestinal system	Loss of appetite, gastric distress

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptom's name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually worsen.

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor with this leaflet, if symptoms do not improve after taking this medicine for about 1 month.
4. If you take this medicine regularly for a long period, consult a physician, pharmacist, or registered distributor.

Dosage and administration

Take the following doses with water or warm water before or between meals.
(Between meals means 2-3 hours after meals.)

Age	One dosage	Daily dosage
15 years and over	1 packet (4.0g)	3 times a day
7 to 14 years old	2/3 packet	
4 to 6 years old	1/2 packet	
2 to 3 years old	1/3 packet	
Less than 2 years old	1/4 packet	

Ingredient and amount

3 packets (4.0 g × 3 : 12.0 g) of this medicine contains 5.6 g of the dried extract (Sokeikakketsuto extract) from the following mixed crude drugs.

Japanese Angelica Root	3 g	Sinomenium Stem and Rhizome	1.5 g
Rehmannia Root	2 g	Notopterygium	1.5 g
Atractylodes Rhizome	2 g	Glehnia Root and Rhizome	1.5 g
Cnidium Rhizome	2 g	Japanese Gentian	1.5 g
Peach Kernel	2 g	Ginger	0.5 g
Poria Sclerotium	2 g	Citrus Unshiu Pee	1.5 g
Peony Root	2.5 g	Angelica Dahurica Root	1.0 g
Achyranthes Root	1.5 g	Glycyrrhiza	1.0 g
Clematis Root	1.5g		

The inactive ingredients contained are Lactose Hydrate, Corn Starch and Magnesium Stearate.

Precautions for storage and handling

- (1) Store in a cool, low humidity place away from direct sunlight.
 - (2) Keep out of reach of children.
 - (3) Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
 - (4) When you take this medicine in divided dose, fold over the opened edge of the packet and store it. The remaining medicine should be taken within 2 days.
- This product is manufactured from natural crude drugs. There may be slight differences in the color and taste of the product. There is no change in the effect.

REQUEST FOR LITERATURE SHOULD BE MADE TO :

Dep. of Medical Information,

Ohsugi Pharmaceutical Co., Ltd.

1-8-6, Yamasaka, Higashisumiyoshi-ku, Osaka 546-0035

Website <https://www.ohsugi-kanpo.co.jp>